

Test na nagibnom stolu u dijagnostici sinkope

The tilt table test in syncope diagnosis

Valentina Sedinić*,
Marina Pavlinić,
Blaženka Marković,
Božica Leško

Klinički bolnički centar Sestre milosrdnice, Zagreb, Hrvatska
University Hospital Centre
"Sestre milosrdnice", Zagreb,
Croatia

KLJUČNE RIJEČI: sinkopa, dijagnostika, sestrinska skrb.

KEYWORDS: syncope, diagnostic, healthcare.

CITATION: *Cardiol Croat.* 2016;11(10-11):566. | DOI: <http://dx.doi.org/10.15836/ccar2016.566>

***ADDRESS FOR CORRESPONDENCE:** Valentina Sedinić, Klinički bolnički centar Sestre milosrdnice, Vinogradska 29, HR-10000 Zagreb, Croatia. / Phone: +385-1-3787-794 / E-mail: valentina.sedinic@kbcsm.hr

ORCID: Valentina Sedinić, <http://orcid.org/0000-0003-0020-7020> • Marina Pavlinić, <http://orcid.org/0000-0001-5734-6342>
Blaženka Marković, <http://orcid.org/0000-0001-6941-8876> • Božica Leško, <http://orcid.org/0000-0001-7423-3999>

Danas je test na nagibnome stolu (eng. *tilt table test*) jedna od nezaobilaznih dijagnostičkih pretraga kod bolesnika sa sinkopom.¹ Nesvjestice, prolazni gubitci svijesti, iznenadna mračenja pred očima, kolapsi i vrtoglavice česte su indikacije za izvođenje navedene pretrage. Sinkope, iako su u osnovi benigne, mogu imati kobne posljedice po bolesnika ako se isti nalazi u prometu ili se ozljedi prilikom pada. Test na nagibnom stolu koristi se primarno za razjašnjenje etiologije sinkope kod anamnestički dvojbenih slučajeva. Ovisno o rezultatu testa (primarno kod kardioinhibitornih sinkopa) može se indicirati ugradnja elektrostimulatora srca.

Prikazom bolesnika prikazat ćemo upotrebu testa na nagibnom stolu u našoj svakodnevnoj kliničkoj praksi.

Today the tilt table test is one of the unavoidable diagnostic tests in patients with syncope.¹ Fainting, transient loss of consciousness, sudden dimming, collapse and dizziness are common indications for performing the test. Syncope, although basically benign, can have fatal consequences for patients if they are engaged in traffic or are injured during the fall. The tilt table test is used primarily to clarify the etiology of syncope in dubious cases. Depending on the result of the test (primarily in cardioinhibitory syncope) it may indicate the implantation of a cardiac pacemaker.

In our case report we will show the use of the tilt table test in our daily clinical practice.

RECEIVED:
September 27, 2016

ACCEPTED:
October 10, 2016



LITERATURE

1. Miller TH, Kruse JE. Evaluation of syncope. *Am Fam Physician.* 2005;72(8):1492-500. PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16273816>